Sign in

Married	***	8	š.
(/\	ogle	COM	nniar
1	///X/#W		
Mille Mill.	Market Barrell	me me s	s area s present

pylori binding GalNac

Search

Advanced Scholar Search Scholar Preferences

Scholar lanytime

include citations

Results 1 - 10 of about 996. (0.04 sec)

PM Simon, PL Goode, A Mobasseri, D Zopf - Infection and immunity, 1997 - iai.highwire.org

- ... Inhibition of Helicobacter pylori Binding to Gastrointestinal Epithelial ... TABLE
- 3. Inhibition of H. pylori binding to HEp-2 monolayers ...

Cited by 151 - Related articles - BL Direct - All 10 versions

Functional adaptation of BabA, the H. pylori ABO blood group antigen binding adhesin- *ns02.info profit

M Aspholm-Hurtig, G Dailide, M Lahmann, A ... - Science, 2004 - sciencemag.org

... group system are formed by terminal N-acetylgalactosamine (GalNAc) or galactose ...

particularly prone to peptic ulcer disease (5). Binding of H. pylori to H1 ...

Cited by 106 - Related articles - All 10 versions

Acidic pH changes receptor **binding** specificity of Helicobacter **pylori**: a binary adhesion ...- **iih.gov port

M Huesca, S Borgia, P Hoffman, CA ... - Infection and immunity, 1996 - Am Soc Microbiol

- ... H. **pylori** to specifically **bind** to the glycolipids gangliotetraosylceramide (Gal
- 1-3GalNac 1-4Gal 1-4Glu cer [Gg 4]) and gangliotriao-sylceramide (GalNac 1 ...

Cited by 89 - Related articles - BL Direct - All 8 versions

- S NORMARK Proc. Natl. Acad. Sci. USA, 1993 National Acad Sciences
- ... Binding is inhibited by neuraminidase and fetuin. The organism also produces sialic

acid-specific hemagglutinins (9). Moreover, H. pylori are able ...

Cited by 174 - Related articles - BL Direct - All 12 versions

Carbohydrate binding specificity of the neutrophil-activating protein of Helicobacter pylori-

- S Teneberg, H Miller-Podraza, HC Lampert, ... Journal of Biological Chemistry, 1997 ASBMB
- ... activating protein from H. **pylori** to glycosphingolipids on ... or abbreviation,
- Glycosphingolipid structure, Binding a, Source. ... 4. Globoside, GalNAc 3Gal 4Gal 4Glc ...

Cited by 62 - Related articles - BL Direct - All 6 versions

- ... 5 Ac alpha 3 Gal is part of the Helicobacter **pylori binding** epitope in polyglycosylceramides ...
- L Johansson, P Johansson, H Miller-Podraza European Journal of Biochemistry, 1999 interscience.wilev.com
- ... nigra, specific for Neu5Ac\(\alpha\)6Gal\(\begin{aligned} \mathbb{GalNAc} \), were used to confirm this distribution.
- H. pylori recognized all three fractions in the binding assay, indicating that ...

Cited by 15 - Related articles - 8L Direct - All 5 versions

The lactosylceramide binding specificity of Helicobacter pylori- *oxfordiournals.org

J Angstrom, S Teneberg, MA Milh, T Larsson, ... - Glycobiology, 1998 - Soc Glycobiology

... H.pylori does bind to gangliotriaosylceramide, but since this binding was concluded to ... presence of an intact acetamido moiety of the terminal GalNAc[beta]4 ...

Cited by 46 - Related articles - 8L Direct - All 4 versions

Helicobacter pylori interactions with human gastric mucin studied with a resonant mirror ...

S Hirmo, E Artursson, G Puu, T Wadström, B ... - Journal of microbiological methods, 1999 - Elsevier ... Displacement of bound H. **pylori** cells from the ... For **binding** competition assays, the bacteria were preincubated ... with HPA, which binds to **GalNAc** residues, was ...

Cited by 22 - Related articles - All 6 versions

... monkey gastric mucins: oligomeric structure, glycoforms and Helicobacter pylori binding.- *** [POF]

S Lindén, T Borén, A Dubois, I Carlstedt - Biochemical Journal, 2004 - pubmedcentral.nih.gov

... with the Elderberry bark (Neu5Ac α 2,6Gal/**GalNAc**) or DBA ... and carbohydrate structure are relevant for H. **pylori** mucin **binding**, statistical correlations ...

Cited by 15 - Related articles - Bt. Direct - All 9 versions

The glycerolipid receptor for Helicobacter pylori (and excenzyme S) is ...- *nih.gov por

CA Lingwood, M Huesca, A Kuksis - Infection and immunity, 1992 - Am Soc Microbiol ... unstable and therefore was not used in these studies. The **binding** of H. **pylori** (13) and of exoenzyme S (12) to receptor lipids was ...

Cited by 93 - Related articles - All 8 versions

	Go	0	0	0	0	0	0	0	0	OS	le	
Result Page:										<u>10</u>		<u>ext</u>

Go to Google Home - About Google - About Google Scholar

©2009 Google